

Algebraic non-hyperbolicity of hyperkähler manifolds with Picard rank greater than one

Ljudmila Kamenova, Misha Verbitsky¹

Abstract. A projective manifold is algebraically hyperbolic if the degree of any curve is bounded from above by its genus times a constant, which is independent from the curve. This is a property which follows from Kobayashi hyperbolicity. We prove that hyperkähler manifolds are not algebraically hyperbolic when the Picard rank is at least 3, or if the Picard rank is 2 and the SYZ conjecture on existence of Lagrangian fibrations is true. We also prove that if the automorphism group of a hyperkähler manifold is infinite then it is algebraically non-hyperbolic.

1 Introduction

In [V1] M. Verbitsky proved that all hyperkähler manifolds are Kobayashi non-hyperbolic. It is interesting to inquire if projective hyperkähler manifolds are also algebraically non-hyperbolic (Definition 2.5). For a given projective manifold algebraic non-hyperbolicity implies Kobayashi non-hyperbolicity. We prove algebraic non-hyperbolicity for projective hyperkähler manifolds with infinite group of automorphisms.

Theorem 1.1: Let M be a projective hyperkähler manifold with infinite automorphism group. Then M is algebraically non-hyperbolic.

If a projective hyperkähler manifold has Picard rank at least three, we show that it is algebraically non-hyperbolic. For the case when the Picard rank equals to two we need an extra assumption in order to prove algebraic non-hyperbolicity. The SYZ conjecture states that a nef parabolic line bundle on a hyperkähler manifold gives rise to a Lagrangian fibration (Conjecture 2.4).

Theorem 1.2: Let M be a projective hyperkähler manifold with Picard rank ρ . Assume that either $\rho > 2$, or $\rho = 2$ and the SYZ conjecture holds. Then M is algebraically non-hyperbolic.

2 Basic notions

Definition 2.1: A *hyperkähler manifold of maximal holonomy* (or *irreducible holomorphic symplectic*) manifold M is a compact complex Kähler manifold with $\pi_1(M) = 0$ and $H^{2,0}(M) = \mathbb{C}\sigma$, where σ is everywhere non-degenerate. From

¹Partially supported by RSCF grant 14-21-00053 within AG Laboratory NRU-HSE.

now on we would tacitly assume that hyperkähler manifolds are of maximal holonomy.

Due to results of Matsushita, holomorphic maps from hyperkähler manifolds are quite restricted.

Theorem 2.2: (Matsushita, [Mat]) Let M be a hyperkähler manifold and $f: M \rightarrow B$ a proper surjective morphism with a smooth base B . Assume that f has connected fibers and $0 < \dim B < \dim M$. Then f is Lagrangian and $\dim_{\mathbb{C}} B = n$, where $\dim_{\mathbb{C}} M = 2n$.

Following Theorem 2.2, we call the surjective morphism $f: M \rightarrow B$ a *Lagrangian fibration* on the hyperkähler manifold M . A dominant map $f: M \dashrightarrow B$ is a *rational Lagrangian fibration* if there exists a birational map $\varphi: M \dashrightarrow M'$ between hyperkähler manifolds such that the composition $f \circ \varphi^{-1}: M' \rightarrow B$ is a Lagrangian fibration. J.-M. Hwang proved that if the base B of a hyperkähler Lagrangian fibration is smooth, then $B \cong \mathbb{P}^n$ (see [Hw]).

Definition 2.3: Given a hyperkähler manifold M , there is a non-degenerate primitive form q on $H^2(M, \mathbb{Z})$, called the *Beauville-Bogomolov-Fujiki form* (or the “*BBF form*” for short), of signature $(3, b_2 - 3)$, and satisfying the *Fujiki relation*

$$\int_M \alpha^{2n} = c \cdot q(\alpha)^n \quad \text{for } \alpha \in H^2(M, \mathbb{Z}),$$

with $c > 0$ a constant depending on the topological type of M . This form generalizes the intersection pairing on K3 surfaces. A detailed description of the form can be found in [Be], [Bog] and [F].

Notice that given a Lagrangian fibration $f: M \rightarrow \mathbb{P}^n$, if h is the hyperplane class on \mathbb{P}^n , and $\alpha = f^*h$, then α belongs to the birational Kähler cone of M and $q(\alpha) = 0$. The following SYZ conjecture states that the converse is also true.

Conjecture 2.4: (Tyurin, Bogomolov, Hassett-Tschinkel, Huybrechts, Sawon) If L is a line bundle on a hyperkähler manifold M with $q(L) = 0$, and such that $c_1(L)$ belongs to the birational Kähler cone of M , then L defines a rational Lagrangian fibration.

For more reference on this conjecture, please see [HT], [Saw], [Hu3] and [V2]. This conjecture is known for deformations of Hilbert schemes of points on K3 surfaces (Bayer–Macrì [BM]; Markman [Mar]), and for deformations of the generalized Kummer varieties $K_n(A)$ (Yoshioka [Y]).

In [V1] M. Verbitsky proved that all hyperkähler manifolds are Kobayashi non-hyperbolic. In [KLV] together with S. Lu we proved that the Kobayashi

pseudometric vanishes identically for K3 surfaces and for hyperkähler manifolds deformation equivalent to Lagrangian fibrations under some mild assumptions. In [De] Demailly introduced the following notion.

Definition 2.5: A projective manifold M is *algebraically hyperbolic* if for any Hermitian metric h on M there exists a constant $A > 0$ such that for any holomorphic map $\varphi: C \rightarrow M$ from a curve of genus g to M we have that $2g - 2 \geq A \int_C \varphi^* \omega_h$, where ω_h is the Kähler form of h .

In this paper all varieties we consider are smooth and projective. For projective varieties, Kobayashi hyperbolicity implies algebraic hyperbolicity ([De]). Here we explore non-hyperbolic properties of projective hyperkähler manifolds. Algebraic non-hyperbolicity implies Kobayashi non-hyperbolicity.

3 Main Results

Proposition 3.1: Let M be a hyperkähler manifold admitting a (rational) Lagrangian fibration. Then M is algebraically non-hyperbolic.

Proof: We use the fact that the fibers of a Lagrangian fibrations are abelian varieties ([Mat]). The isogeny self-maps on an abelian variety provide curves of fixed genus and arbitrary large degrees, and therefore they are algebraically non-hyperbolic.

An alternative way of proving this proposition is by using the following result whose proof was suggested by Prof. K. Oguiso.

Lemma 3.2: If a hyperkähler manifold M admits a Lagrangian fibration, then there exists a rational curve on M .

Indeed, in [HO] J.-M. Hwang and K. Oguiso give a Kodaira-type classification of the general singular fibers of a holomorphic Lagrangian fibration. All of the general singular fibers are covered by rational curves. The locus of singular fibers is non-empty (e.g., Proposition 4.1 in [Hw]), and therefore there is a rational curve on M .

According to Lemma 3.2, M contains a rational curve, and therefore, M is algebraically non-hyperbolic. This finishes the proof of Proposition 3.1. ■

Lemma 3.3: Let M be a projective hyperkähler manifold with infinite automorphism group Γ . Consider the natural map $f: \Gamma \rightarrow \text{Aut}(H^{1,1}(M))$. Then the elements of the Kähler cone have infinite orbits with respect to $f(\Gamma)$.

Proof: See the discussion in section 2 of [O2]. ■

Lemma 3.4: Let M be a projective hyperkähler manifold, and Γ its automorphism group. Consider the natural map $g : \Gamma \longrightarrow \text{Aut}(H_{tr}^2(M)) \times \text{Aut}(H^{1,1}(M))$. Then $g(\Gamma)$ is finite in the first component $\text{Aut}(H_{tr}^2(M))$.

Proof: This has been proven by Oguiso, see [O1]. The idea is that the BBF form restricted to the transcendental part $H_{tr}^2(M)$ is of K3-type. Then we can apply Zarhin's theorem (Theorem 1.1.1 in [Z]) to deduce that $g(\Gamma) \subset \text{Aut}(H_{tr}^2(M))$ is finite. ■

Theorem 3.5: Let M be a projective hyperkähler manifold with infinite automorphism group. Then M is algebraically non-hyperbolic.

Proof: In the notations introduced above, for any Kähler class w on M , its $f(\Gamma)$ -orbit is infinite by Lemma 3.3. Fix a polarization w on M with normalization $q(w) = 1$. For a given constant $C > 0$ consider the set

$$\mathcal{D}_C = \{x \in H^{1,1}(M, \mathbb{Z}) \mid q(x) \geq 0, \quad q(x, w) \leq C\}.$$

Notice that \mathcal{D}_C is compact. Indeed, $y = x - q(x, w)w$ is orthogonal to w with respect to the BBF form q . The quadratic form q is of type $(1, \rho - 1)$ on $H^{1,1}(M, \mathbb{Z})$ and since $q(w) > 0$, the restriction $q|_{w^\perp}$ is negative-definite. A direct computation shows that $q(y) = q(x) - 2q(x, w)^2 + q(x, w)^2 q(w) = q(x) - q(x, w)^2 \geq -C^2$. The set \mathcal{D}_C is equivalent to the set of elements $\{y \in w^\perp \mid q(y) \geq -C^2\}$, which is compact because $q|_{w^\perp}$ is negative-definite. Since the set \mathcal{D}_C is compact, $\sup_{x \in \Gamma \cdot \eta} \deg x = \infty$, which means there is a class of a curve η with $q(\eta) > 0$. However, all curves in the orbit $\Gamma \cdot \eta$ have constant genus. Since their degrees could be arbitrarily high, then M is algebraically non-hyperbolic. ■

Lemma 3.6: Let M be a hyperkähler manifold such that the positive cone does not coincide with the Kähler cone. Then M contains a rational curve.

Proof: This is a classical result that Boucksom and Huybrechts knew in the early 2000's [Bou, Hu2]. ■

Theorem 3.7: Let M be a hyperkähler manifold with Picard rank ρ . Assume that either $\rho > 2$ or $\rho = 2$ and the SYZ conjecture holds. Then M is algebraically non-hyperbolic.

Proof: Notice that the Hodge lattice $H^{1,1}(M, \mathbb{Z})$ of a hyperkähler manifold has signature $(1, k)$. Therefore, for $\rho \geq 2$, the Hodge lattice contains a vector with positive square, and M is projective ([Hu1]). First, consider the case when $\rho > 2$. If the Kähler cone coincides with the positive cone, then the automorphism group $\text{Aut}(M)$ is commensurable with the group of isometries $SO(H^2(M, \mathbb{Z}))$ (Theorem 2.17 in [AV]) preserving the Hodge type. By Lemma 3.4, this group

is commensurable with the group of isometries of the Hodge lattice $H^{1,1}(M, \mathbb{Z})$. By Borel and Harish-Chandra's theorem ([BHC]), if $\rho > 2$, any arithmetic subgroup of $SO(1, \rho - 1)$ is a lattice. However, Borel density theorem implies that any lattice in a non-compact simple Lie group is Zariski dense ([Bor]). Therefore, for $\rho > 2$, $SO(H^{1,1}(M, \mathbb{Z}))$ is infinite. In this case $\text{Aut}(M)$ is also infinite and we can apply Theorem 3.5. On the other hand, if the Kähler cone does not coincide with the positive cone, then by Lemma 3.6 there is a rational curve on M . Therefore, M is algebraically non-hyperbolic.

Now let $\rho = 2$. Assume the positive cone and the Kähler cone coincide. If there is no $\eta \in H^{1,1}(M, \mathbb{Z})$ with $q(\eta) = 0$, then by Theorem 87 in [Di], $SO(H^{1,1}(M, \mathbb{Z}))$ is isomorphic to $\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. Therefore, both $SO(H^{1,1}(M, \mathbb{Z}))$ and $\text{Aut}(M)$ are infinite and we can apply Theorem 3.5. If there is $\eta \in H^{1,1}(M, \mathbb{Z})$ with $q(\eta) = 0$, then the SYZ conjecture implies that η defines a rational fibration on M and we could apply Proposition 3.1. If $\rho = 2$ and the positive and the Kähler cones are different (i.e., the positive cone is divided into Kähler chambers), then there is a nef class $\eta \in H^{1,1}(M, \mathbb{Z})$ with $q(\eta) = 0$. Since we assumed that the SYZ conjecture holds, the class η defines a Lagrangian fibration on M . Applying Proposition 3.1 we conclude that M is algebraically non-hyperbolic. ■

Remark 3.8: We conjecture that all projective hyperkähler manifolds are algebraically non-hyperbolic. However, our proof fails for manifolds with Picard rank 1.

Acknowledgments. This work was inspired by a question of Erwan Rousseau about algebraic non-hyperbolicity of hyperkähler manifolds. The paper was started while the first-named author was visiting Université libre de Bruxelles and she is grateful to Joel Fine for his hospitality. It was finished at the SCGP during the second-named author's stay there. We are grateful to the SCGP for making this possible.

References

- [AV] Amerik, E., Verbitsky, M., *Construction of automorphisms of hyperkähler manifolds*, arXiv:1604.03079 [math.AG].
- [BM] Bayer, A., Macri, E., *MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations*, Invent. Math. **198** (2014) 505 - 590.
- [Be] Beauville, A., *Variétés Kähleriennes dont la première classe de Chern est nulle*. J. Diff. Geom. **18** (1983) 755 - 782.
- [Bog] Bogomolov, F., *Hamiltonian Kähler manifolds*, Sov. Math. Dokl. **19** (1978) 1462 - 1465.

- [Bor] A. Borel, *Density properties for certain subgroups of semi-simple groups without compact components*, Ann. of Math. (2) **72** (1960), 179–188.
- [Bou] Boucksom, S., *Le cône kählérien d’une variété hyperkählérienne*, C. R. Acad. Sci. Paris Ser. I Math. **333** (2001), no. 10, 935–938.
- [BHC] Borel, A., Harish-Chandra, *Arithmetic subgroups of algebraic groups*, Ann. Math. (2) **75** (1962) 485 - 535.
- [CR] Curtis, W. C., Reiner, I., *Representation theory of finite groups and associative algebras*, Interscience, New York (1962).
- [De] Demailly, J.-P., *Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials*, Proc. Symp. Pure Math., Vol **62.2** (1997) 285 - 360.
- [Di] Dickson, L. E., *Introduction to the theory of numbers*, Dover Publ. Inc., New York (1954).
- [F] Fujiki, A., *On the de Rham Cohomology Group of a Compact Kähler Symplectic Manifold*, Adv. Stud. Pure Math. **10** (1987) 105 - 165.
- [HT] Brendan Hassett, Yuri Tschinkel, *Rational curves on holomorphic symplectic fourfolds*, arXiv:math/9910021, Geom. Funct. Anal. **11** (2001), no. 6, 1201–1228.
- [Hu1] Huybrechts, D., *Compact hyper-Kähler manifolds: basic results*, Invent. Math. **135** (1999) 63 - 113.
- [Hu2] Huybrechts, D., *The Kähler cone of a compact hyperkähler manifold*, Math. Ann. **326** (2003), no. 3, 499–513, arXiv:math/9909109.
- [Hu3] Huybrechts, Daniel, *Compact hyperkähler manifolds, Calabi-Yau manifolds and related geometries*, Universitext, Springer-Verlag, Berlin, 2003, Lectures from the Summer School held in Nordfjordeid, June 2001, pp. 161–225.
- [Hw] Hwang, J.-M., *Base manifolds for fibrations of projective irreducible symplectic manifolds*, Invent. Math. **174** (2008) 625 - 644.
- [HO] Hwang, J.-M., Oguiso, K., *Characteristic foliation on the discriminant hypersurface of a holomorphic Lagrangian fibration*, Am. J. Math. **131** (2009) 981 - 1007.
- [KLV] Kamenova, L., Lu, S., Verbitsky, M., *Kobayashi pseudometric on hyperkahler manifolds*, J. London Math. Soc. (2014) **90** (2): 436 - 450.
- [Mar] Markman, E., *Lagrangian fibrations of holomorphic-symplectic varieties of $K3^{[n]}$ -type*, Springer Proceedings in Mathematics & Statistics **71** (2014) 241 - 283. Algebraic and complex geometry. In honour of Klaus Hulek’s 60th birthday.

- [Mat] Matsushita, D., *On fibre space structures of a projective irreducible symplectic manifold*, Topology **38** (1999), No. 1, 79 - 83. Addendum, Topology **40** (2001) No. 2, 431 - 432.
- [O1] Oguiso, K., *Automorphisms of hyperähler manifolds in the view of topological entropy*, AMS, Contemp. Math. 422 (2007) 173 - 185.
- [O2] Oguiso, K., *Bimeromorphic automorphism groups of non-projective hyperkähler manifolds - a note inspired by C. T. McMullen*, J. Differ. Geom. 78 (2008) 163 - 191.
- [Saw] J. Sawon, *Abelian fibred holomorphic symplectic manifolds*, Turkish Jour. Math. 27 (2003), no. 1, 197-230.
- [V1] Verbitsky, M., *Ergodic complex structures on hyperkähler manifolds*, Acta Math. **215** (2015) 161 - 182.
- [V2] Verbitsky, M., *Hyperkahler SYZ conjecture and semipositive line bundles*, arXiv:0811.0639, GAFA 19, No. 5 (2010) 1481-1493.
- [Y] Yoshioka, K., *Bridgeland's stability and the positive cone of the moduli spaces of stable objects on an abelian surface*, arXiv:1206.4838 [math.AG].
- [Z] Zarhin, Yu., *Hodge groups of K3 surfaces*, J. Reine Angew. Math. 341 (1983) 193 - 220.

LJUDMILA KAMENOVA
 DEPARTMENT OF MATHEMATICS, 3-115
 STONY BROOK UNIVERSITY
 STONY BROOK, NY 11794-3651, USA,
 kamenova@math.sunysb.edu

MISHA VERBITSKY
 LABORATORY OF ALGEBRAIC GEOMETRY,
 NATIONAL RESEARCH UNIVERSITY HSE,
 FACULTY OF MATHEMATICS, 7 VAVILOVA STR.,
 MOSCOW, RUSSIAN FEDERATION,
 verbit@mccme.ru, also:
 UNIVERSITÉ LIBRE DE BRUXELLES, CP 218,
 BD DU TRIOMPHE, 1050 BRUSSELS, BELGIUM